# HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated SUPERVALU INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## Equaline 44-334B

# Active ingredients (in each caplet)

Acetaminophen 250 mg Aspirin 250 mg (NSAID)\* Caffeine 65 mg \*nonsteroidal anti-inflammatory drug

## **Purpose**

Pain reliever Pain reliever Pain reliever aid

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - arthritis
  - a cold
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

• more than 4,000 mg of acetaminophen in 24 hours

- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

#### Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

# Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

## Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur

These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

## Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

# **Inactive ingredients**

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

#### Questions or comments?

1-855-423-2630

### Principal display panel

**EQUALINE®** 

compare to Excedrin® Extra Strength active ingredients\*\*

NDC 41163-334-12

extra strength

headache relief acetaminophen aspirin *(NSAID)* and caffeine

pain reliever/pain reliever aid

100 caplets

actual size

#### **DOES NOT CONTAIN GLUTEN**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

### 50844 REV1119B33412

\*\*This product is not manufactured or distributed by GSK Consumer Healthcare S.A., owner of the registered trademark Excedrin® Extra Strength.

# DISTRIBUTED BY UNFI PROVIDENCE, RI 02908 USA

855-423-2630



Equaline 44-334B

acetaminophen, aspirin, caffeine tablet, film coated

<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-334
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	250 mg		
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg		
CAFFEINE (UNII: 3G6 A5W338E) (CAFFEINE - UNII:3G6 A5W338E)	CAFFEINE	65 mg		

Inactive Ingredients				
Ingredient Name	Strength			
STARCH, CORN (UNII: O8232NY3SJ)				
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)				
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SODIUM LAURYL SULFATE (UNII: 368 GB5141J)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)				

Product Characteristics				
Color	WHITE	Score	no score	
Shape	OVAL	Size	17mm	
Flavor		Imprint Code	44;334	
Contains				

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-334- 12	1 in 1 CARTON	11/0 1/20 17	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:41163-334- 08	1 in 1 CARTON	11/0 1/20 17	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FINAL	part343	11/0 1/20 17		

# Labeler - SUPERVALU INC. (006961411)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(41163-334)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		038154464	PACK(41163-334)	

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		868734088	PACK(41163-334)	

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		967626305	PACK(41163-334)

Revised: 4/2020 SUPERVALU INC.